arXiv:1406.4847v1 [math.NT] 18 Jun 2014

A NOTE REGARDING PERMUTATION BINOMIALS OVER $\mathbb{F}_{q^2}$

STEPHEN D. LAPPANO

Abstract. Let $f = ax + x^{r(q-1)+1} \in \mathbb{F}_{q^2}^*[x], r \in \{5, 7\}$. We give explicit conditions on the values (q, a) for which f is a permutation polynomials of $\mathbb{F}_{q^2}$.

1. Introduction

A polynomial $f \in \mathbb{F}_q[x]$ is called a *permutation polynomial* (PP) if the mapping $\mathtt{x} \mapsto f(\mathtt{x})$ induces a permutation on $\mathbb{F}_q$. Permutation polynomials taking simple algebraic forms are particularly interesting. While permutation monomials are quite obvious, the situation for permutation binomials is not as well understood. As a result much research has been concerned with finding, classifying and understanding permutation binomials. The main results of the present paper are the following theorems.

Theorem 1.1. *Let $f(x) = ax + x^{5q-4} \in \mathbb{F}_{q^2}[x]$. Then f is a PP of $\mathbb{F}_{q^2}$ iff one of the following occurs:*

1. *$q = 2^{4k+2}$ and $a^{\frac{q+1}{5}} \neq 1$ is a fifth root of unity*
2. *$q = 3^2$ and a^2 is a root of $(1+x)(1+x^2)(2+x+x^2)(2+2x+x^2)(1+x+x^2+x^4)(1+x^2+x^3+x^4)(1+2x+x^2+2x^3+x^4)$*
3. *$q = 19$ and a^4 is a root of $(1+x)(2+x)(3+x)(4+x)(5+x)(9+x)(10+x)(13+x)(17+x)(16+3x+x^2)(1+4x+x^2)(6+18x+x^2)$*
4. *$q = 29$ and $a^6 \in \{15, 18, 22, 23\}$*
5. *$q = 7^2$ and a^{10} is a root of $(1+4x+x^2)$*
6. *$q = 59$ and a^{12} is a root of $(4+x)(55+x)(x^2+36)$*
7. *$q = 2^6$ and a^{13} is a root of $(1+x+x^2)(1+x+x^3)$*

Theorem 1.2. *Let $f(x) = ax + x^{7q-6} \in \mathbb{F}_{q^2}[x]$. Then f is a PP of $\mathbb{F}_{q^2}$ iff one of the following occurs:*

1. *$q = 13$ and a^2 is a root of $(1+x)(2+x)(3+x)(4+x)(5+x)(6+x)(7+x)(8+x)(9+x)(10+x)(11+x)(12+x+x^2)(9+2x+x^2)(10+3x+x^2)(9+4x+x^2)(12+4x+x^2)(10+5x+x^2)(3+6x+x^2)(1+7x+x^2)(4+7x+x^2)(1+8x+x^2)(12+9x+x^2)(1+10x+x^2)(3+12x+x^2)(4+12x+x^2)(12+12x+x^2)$*
2. *$q = 3^3$ and a^4 is a root of $(2+x+x^2+x^3)(1+2x+x^2+x^3)(1+x+2x^2+x^3)(2+2x+2x^2+x^3)(1+2x+x^2+2x^3+x^4+2x^5+x^6)$*
3. *$q = 41$ and a^6 is a root of $(9+x)(10+x)(26+x)(30+x)(32+x)(34+x)(35+x)(37+x)(39+2x+x^2)(1+14x+x^2)(20+40x+x^2)$*

It is worth noting the polynomial f in Theorems 1.1 and 1.2 can be written as $f = xh(x^{q-1})$, where $h(x) = a + x^r$. Zieve [5] considered polynomials of this type and under the assumption $a \in \mathbb{F}_{q^2}$ is a (q+1)st root of unity Theorem 1.1 (1) follows from [5,Cor 5.3]. The approach of this paper is quite different from that of [5], as

no restrictions are placed on the nature of $a \in \mathbb{F}_{q^2}$. We employ a method similar to that of [1, 2] and expand the sum $\sum_{x\in\mathbb{F}_{q^2}} f(x)^s$. Surprisingly, considering only a few values of s give explicit results. In [3] permutation binomials of the form $f = ax + x^{3q-2}$ were completely characterized. The present paper extends this result to polynomials of the form $f = ax + x^{5q-4}$ and $f = ax + x^{7q-6}$. We conclude with a conjecture regarding the behavior of $f = ax + x^{r(q-1)+1}$ for odd primes r.

2. Computations

Let $f = ax + x^{5q-4} \in \mathbb{F}_{q^2}^*[x]$ and let $0 \le \alpha, \beta \le q-1$. We compute

$$\begin{aligned}\sum_{x\in\mathbb{F}_{q^2}} f(x)^{\alpha+\beta q} &= \sum_{x\in\mathbb{F}_{q^2}} (ax + x^{5q-4})^\alpha (a^q x^q + x^{5-4q})^\beta \\ &= \sum_{x\in\mathbb{F}_{q^2}} \sum_{i,j} \binom{\alpha}{i} (ax)^{\alpha-i} x^{(5q-4)i} \binom{\beta}{j} (a^q x^q)^{(\beta-j)} x^{(5-4q)j} \\ &= a^{\alpha+\beta q} \sum_{i,j} \binom{\alpha}{i}\binom{\beta}{j} a^{-i-jq} \sum_{x\in\mathbb{F}_{q^2}} x^{\alpha+\beta q+5(q-1)(i-j)}. \qquad (2.1)\end{aligned}$$

It is clear the inner sum is 0 unless $\alpha + \beta q \equiv (\bmod q-1)$, thus $\alpha + \beta = q-1$.

With $0 \le \alpha \le q-1$ and $\beta = q-1-\alpha$ (2.1) becomes

$$-a^{(\alpha+1)(1-q)} \sum_{-\alpha-1+5(i-j)\equiv 0 (\mathrm{mod} q+1)} \binom{\alpha}{i}\binom{q-1-\alpha}{j} a^{-i-jq} \qquad (2.2)$$

Since $0 \le i \le q-1$ and $0 \le \beta \le q-1-\alpha$ it follows

$$4\alpha + 4 - 5q \le -\alpha - 1 + 5(i-j) \le 4\alpha - 1 \qquad (2.3)$$

Define $\Gamma(q,\alpha) := \{n \in (q+1)\mathbb{Z} : 4\alpha + 4 - 5q \le n \le 4\alpha - 1\}$. Now we have

$$\sum_{x\in\mathbb{F}_{q^2}} f(x)^{\alpha+\beta q} = -a^{(\alpha+1)(1-q)} \Lambda(q,\alpha,a) \qquad (2.4)$$

where

$$\Lambda(q,\alpha,a) = \sum_{-\alpha-1+5(i-j)\in\Gamma(q,\alpha)} \binom{\alpha}{i}\binom{q-1-\alpha}{j} a^{-i-jq} \qquad (2.5)$$

Now Hermite's criterion implies f is a PP of $\mathbb{F}_{q^2}$ if and only if 0 is the only root of f in $\mathbb{F}_{q^2}$ and

$$\Lambda(q,\alpha,a) = 0 \text{ for each } 0 \le \alpha \le q-1. \qquad (2.6)$$

Remark 2.1. *Notice if $q+1 \equiv 0 \pmod 5$, then 0 is the only root of f if and only if $a^{\frac{q+1}{5}} \neq 1$.*

Lemma 2.2. *If f is a PP of $\mathbb{F}_{q^2}$, then $q+1 \equiv 0$ mod 5.*

Proof. Assume f is a PP and $q \ge 5$. First suppose $5 \le q < 8$. Note that

$$\Gamma(q,0) = \{-3(q+1), -2(q+1), -(q+1)\}.$$

By (2.6) we have

$$0 = \Lambda(5,0,a) = \binom{5-1}{1} a^{-5} = -a^{-5}$$

and

$$0 = \Lambda(7, 0, a) = \binom{7-1}{1} a^{-21} = -a^{-21}.$$

In either case we have a contradiction. Now suppose $q \geq 8$. In this case notice

$$\Gamma(q, 0) = \{-4(q+1), -3(q+1), -2(q+1), -(q+1)\}.$$

So again by (2.6) we have

$$0 = \Lambda(q, 0, a) = \sum_{k=1}^{4} \binom{q-1}{\frac{k(q+1)-1}{5}}^{*} a^{-(\frac{k(q+1)-1}{5})q}$$

where

$$\binom{n}{m}^{*} = \begin{cases} \binom{n}{m} & \text{if } m \in \mathbb{Z} \\ 0 & \text{otherwise} \end{cases}.$$

Now if $(q+1) \not\equiv 0 \pmod 5$ then exactly one of $\binom{q-1}{\frac{k(q+1)-1}{5}}^{*}$ is nonzero which contradicts (2.6). Thus if f is a PP of $\mathbb{F}_{q^2}$ we must have $q+1 \equiv 0 \pmod 5$. □

Remark 2.3. *Assume $q+1 \equiv 0 \pmod 5$ and $\alpha > 0$. The previous lemma together with (2.5) imply the sum $\Lambda(q, \alpha, 0)$ is empty unless $\alpha + 1 \equiv 0 \pmod 5$.*

Lemma 2.4. *Assume $q+1 \equiv 0 \pmod 5$, $\alpha > 0$, $\alpha+1 \equiv 0 \pmod 5$ and $q \geq 4\alpha+8$. Set $v = a^{-\frac{q+1}{5}}$, then*

$$\Lambda(q, \alpha, a) = (-a)^{\frac{\alpha+1}{5}q} \sum_{i=0}^{\alpha} (-1)^i \binom{\alpha}{i} \sum_{l=0}^{4} \binom{i + \frac{4\alpha-1+l}{5}}{\alpha} v^{5i+l}.$$

Proof. Since $q \geq 4\alpha + 8$ we have

$$\Gamma(q, \alpha) = \{-4(q+1), -3(q+1), -2(q+1), -(q+1), 0\}.$$

Using (2.5) we see

$$\begin{aligned}
\Lambda(q, \alpha, a) &= \sum_{-\alpha-1+5(i-j) \in \Gamma(q,\alpha)} \binom{\alpha}{i} \binom{q-1-\alpha}{j} a^{-i-jq} \\
&= \sum_{i=0}^{\alpha} \binom{\alpha}{i} \sum_{l=0}^{4} \binom{-1-\alpha}{\frac{1}{5}(l(q+1)-\alpha-1)+i} a^{-i-[\frac{1}{5}(l(q+1)-\alpha-1)+i]q} \\
&= (-a)^{\frac{q+1}{5}q} \sum_{i=0}^{\alpha} \binom{\alpha}{i} \sum_{l=0}^{4} (-1)^i \binom{\frac{1}{5}(l(q+1)-\alpha-1)+i+\alpha}{\alpha} a^{-\frac{q+1}{5}(l+5i)} \\
&= (-a)^{\frac{\alpha+1}{5}q} \sum_{i=0}^{\alpha} (-1)^i \binom{\alpha}{i} \sum_{l=0}^{4} \binom{i + \frac{4\alpha-1+l}{5}}{\alpha} v^{5i+l}.
\end{aligned}$$

Between the second and third line we use $\binom{-m}{n} = (-1)^n \binom{n+m-1}{m-1}$. □

Lemma 2.5. *Assume $q+1 \equiv 0 \pmod 5$ and $\alpha > 0$. $\Gamma(q, \alpha)$ contains exactly five consecutive multiples of $q+1$ unless $\alpha = \frac{q-1}{2} \in \mathbb{Z}$ or $\alpha = \frac{q-3}{4} \in \mathbb{Z}$.*

Proof. We may assume $\alpha+1 \equiv 0 \pmod 5$. Since $\Gamma(q,\alpha)$ in contained in the interval $[4\alpha+4-5q, 4\alpha-1]$ which has length $5(q-1)$ we must have $4 \leq |\Gamma(q,\alpha)| \leq 5$.

Suppose $4 = |\Gamma(q,\alpha)|$. Choose k so $\{(k-3)(q+1),(k-2)(q+1),(k-1)(q+1),k(q+1)\} = \Gamma(q,\alpha)$. Note $q \geq 6$ and $\alpha > 0$ force $k \in \{0,1,2,3\}$. We have the following inequalities

$$\begin{cases} 4\alpha - 1 \leq k(q+1) + q \\ (k-3)(q+1) - q \leq 4\alpha + 4 - 5q. \end{cases} \tag{2.7}$$

Since $4\alpha - 1, 4\alpha + 4 - 5q \equiv 0 \pmod 5$ it follows

$$\begin{cases} 4\alpha - 1 \leq k(q+1) + q - 4 \\ (k-3)(q+1) - q + 4 \leq 4\alpha + 4 - 5q. \end{cases} \tag{2.8}$$

Since $a \leq q-1$ and $0 \leq k \leq 3$, taking the difference of the inequalities in (2.8) reveals $k = 0$ which gives $\alpha = \frac{q-3}{4}$ or $k = 1$ which gives $\alpha = \frac{q-1}{2}$. □

Lemma 2.6. *Assume $q+1 \equiv 0 \pmod 5$ and $y := a^{\frac{q+1}{5}} \neq 1$ is a 5th root of unity. Then for $1 \leq \alpha \leq q-1$ we have*

$$\Lambda(q,\alpha,a) = \begin{cases} -a^{-\frac{1}{5}(\alpha+1)}(y^{-1}+1+y+y^2) & \text{if } \alpha = \frac{q-1}{2}, \ \ \alpha \in \mathbb{Z} \\ -a^{-\frac{1}{5}(\alpha+1)}(1+y+y^2+y^3) & \text{if } \alpha = \frac{q-3}{4}, \ \ \alpha \in \mathbb{Z} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We may assume $\alpha + 1 \equiv 0 \pmod 5$. First suppose $\alpha \neq \frac{q-3}{4}, \frac{q-1}{2}$. Let K be a set of five consecutive integers such that $K(q+1) = \Gamma(q,\alpha)$. Now we have

$$\begin{aligned} \Lambda(q,\alpha,a) &= \sum_{-\alpha-1+5(i-j)\in\Gamma(q,\alpha)} \binom{\alpha}{i}\binom{q-1-\alpha}{j} a^{-i-jq} \\ &= \sum_{k\in K} \sum_{-\alpha-1+5(i-j)=k(q+1)} \binom{\alpha}{i}\binom{q-1-\alpha}{j} a^{-i+j} \\ &= \sum_{k\in K} a^{-\frac{1}{5}[\alpha+1+k(q+1)]} \sum_{i-j=\frac{1}{5}[\alpha+1+k(q+1)]} \binom{\alpha}{\alpha-i}\binom{q-1-\alpha}{j} \\ &= a^{-\frac{1}{5}(\alpha+1)} \sum_{k\in K} y^{-k} \sum_{\alpha-i+j=\frac{1}{5}[4\alpha-1-k(q+1)]} \binom{\alpha}{\alpha-i}\binom{q-1-\alpha}{j} \\ &= a^{-\frac{1}{5}(\alpha+1)} \sum_{k\in K} y^{-k} \binom{q-1}{\frac{1}{5}[4\alpha-1-k(q+1)]} \\ &= -a^{-\frac{1}{5}(\alpha+1)} \sum_{k\in K} y^{-k} = 0. \end{aligned} \tag{2.9}$$

Now suppose $\alpha = \frac{q-1}{2}$. By the above computation and the previous lemma we have $K = \{-2,-1,0,1\}$, thus

$$\Lambda(q,\alpha,a) = -a^{-\frac{1}{5}(\alpha+1)}(y^{-1}+1+y+y^2). \tag{2.10}$$

Similarly if $\alpha = \frac{q-3}{4}$, we have

$$\Lambda(q,\alpha,a) = -a^{-\frac{1}{5}(\alpha+1)}(1+y+y^2+y^3). \tag{2.11}$$

□

3. Proof of the Theorem

Proof. (Theorem1.1)
($\Leftarrow$) Cases (ii)-(vii) are easily verified by a computer. Assume (i), that is $q = 2^{4k+2}$ and $a^{\frac{q+1}{5}} \neq 1$ is a fifth root of unity. Lemma 2.6 gives $\Lambda(q, \alpha, a) = 0$ for each $0 \leq \alpha \leq q-1$ and 0 is the only root of f, so f is a PP by (2.6)

($\Rightarrow$) Assume f is a PP. By Lemma (2.2) we have $q+1 \equiv 0 \pmod 5$. Let $y := a^{\frac{q+1}{5}}$. If $y \neq 1$ is a fifth root of unity then Lemma 2.6 implies q must be even. Thus $q = 2^{4k+2}$ and we have case (i).

Now suppose $1 + y + y^2 + y^3 + y^4 \neq 0$. The sum in the RHS of Lemma 2.4 is a polynomial in $v(= y^{-1})$ and can be easily computed for small values of α with the help of a computer algebra system. For a few values of α we find

$$\Lambda(q, \alpha, a) = (-a)^{\frac{\alpha+1}{5}} v(1+v+v^2+v^3+v^4) \begin{cases} 5^{-4} g_4(v) & \text{if } \alpha = 4, q \geq 24 \\ 5^{-10} g_9(v) & \text{if } \alpha = 9, q \geq 44 \\ 5^{-16} g_{14}(v) & \text{if } \alpha = 14, q \geq 64 \\ 5^{-28} g_{24}(v) & \text{if } \alpha = 24, q \geq 104. \end{cases} \tag{3.1}$$

The polynomials g_α are given in the appendix. Write $R(p_1, p_2)$ for the resultant of polynomials p_1, p_2. Then

$$GCD(R(g_4, g_9), R(g_4, g_{14})) = 2^{15} 3^3 5^{197}. \tag{3.2}$$

Thus if $q \geq 64$ we must have $p\ (= char\mathbb{F}_{q^2}) \in \{2, 3\}$. Since $q+1 \equiv 0 \pmod 5$ there are only a few prime powers $q < 64$ with $p\ (= char\mathbb{F}_{q^2}) \neq 2, 3$.

- When $q = 19$, a computer search results in case (iii)
- When $q = 29$, a computer search results in case (iv)
- When $q = 49$, a computer search results in case (v)
- When $q = 59$, a computer search results in case (vi)

When $p = 2$ we have $GCD(g_4, g_{24}) = x$ thus $q < 104$. Since $q + 1 \equiv 0 \pmod 5$ and $q > 4$ we only need to consider $q = 64$. A computer search results in case (vii).

When $p = 3$ we have $GCD(g_4, g_9) = 1$ thus $q < 44$. Again since $q + 1 \equiv 0 \pmod 5$ we only need to consider $q = 9$. A computer search results in case (ii). □

Theorem 1.2 is proved using a similar method which leads us to the following conjecture.

Conjecture 3.1. *Let $r > 2$ be a fixed prime. If both $(q+1) \equiv 0 \pmod r$ and $a^{\frac{q+1}{r}}$ is not an $r-th$ root of unity; we conjecture there are only finitely many values (q, a) for which $f = ax + x^{r(q-1)+1} \in \mathbb{F}_{q^2}^*[x]$ is a permutation polynomial of $\mathbb{F}_{q^2}$.*

4. Appendix

$$\begin{aligned} g_4(x) = g_4(x) = 44 &+ 75x + 115x^2 + 165x^3 - 2899x^4 - 1152x^5 - 1465x^6 - 1825x^7 \\ &- 2235x^8 + 25427x^9 + 4122x^{10} + 4805x^{11} + 5555x^{12} + 6375x^{13} - 58357x^{14} \\ &- 3514x^{15} - 3915x^{16} - 4345x^{17} - 4805x^{18} + 38454x^{19}. \end{aligned}$$

$$\begin{aligned}g_9(x) &= 35464 + 47564x + 41954x^2 - 3211x^3 - 121771x^4 + 3307018x^5 + 7237538x^6 \\ &+ 13764443x^7 + 24034088x^8 - 399905587x^9 - 247096418x^{10} - 374180158x^{11} \\ &- 550629013x^{12} - 791033308x^{13} + 10166063897x^{14} + 3488769646x^{15} \\ &+ 4732145066x^{16} + 6330108851x^{17} + 8362735316x^{18} - 90589540129x^{19} \\ &- 20005926320x^{20} - 25545256270x^{21} - 32342528845x^{22} - 40628485270x^{23} \\ &+ 389225321705x^{24} + 56878393066x^{25} + 69794856416x^{26} + 85151720951x^{27} \\ &+ 103326084566x^{28} - 901674492499x^{29} - 84876624338x^{30} - 101203850008x^{31} \\ &- 120170621113x^{32} - 142129134058x^{33} + 1152208354517x^{34} + 63621134698x^{35} \\ &+ 74215593188x^{36} + 86304925343x^{37} + 100064515838x^{38} - 764098747192x^{39} \\ &- 18879570941x^{40} - 21644171461x^{41} - 24754360696x^{42} - 28246292086x^{43} \\ &+ 205243145184x^{44}.\end{aligned}$$

$$\begin{aligned}g_{14}(x) &= 35781192 + 45103176x + 34642700x^2 - 11075651x^3 - 104451417x^4 + 3431018744x^5 \\ &+ 5425819552x^6 + 5815099340x^7 + 1030592053x^8 - 15702529689x^9 + 367570246196x^{10} \\ &+ 915262570648x^{11} + 1948805856320x^{12} + 3779801961517x^{13} - 625534328221 81x^{14} \\ &- 47505978162672x^{15} - 79121737743896x^{16} - 127716748824160x^{17} - 200786722216499x^{18} \\ &+ 2746238364681602x^{19} + 1263694841543105x^{20} + 1868009627371040x^{21} \\ &+ 2718322826664950x^{22} + 3900065855839735x^{23} - 46407807995168830x^{24} \\ &- 15130762640643679x^{25} - 20922156001611712x^{26} - 28643147196518350x^{27} \\ &- 38852104359262613x^{28} + 415138746369911354x^{29} + 101011800565346642x^{30} \\ &+ 1335461812744641 11x^{31} + 1753166346556279 70x^{32} + 228624694439538979x^{33} \\ &- 224096513124747 7702x^{34} - 414819831279200662x^{35} - 530625761486025481x^{36} \\ &- 675157922777947390x^{37} - 854715990669328349x^{38} + 7803518358751564382x^{39} \\ &+ 1100711546227581974x^{40} + 137238807860242 3607x^{41} + 170398886116303 9670x^{42} \\ &+ 2107260628357829383x^{43} - 181247910088821 24634x^{44} - 192372297288913 2730x^{45} \\ &- 234956665678498 6915x^{46} - 286002358515015 0200x^{47} - 347009611360053 9985x^{48} \\ &+ 283640721702216 84830x^{49} + 220001557207349 5862x^{50} + 264158510438503 8536x^{51} \\ &+ 316299048136301 8525x^{52} + 377715509203952 9639x^{53} - 295424090916579 57562x^{54} \\ &- 158405377176426 2386x^{55} - 187482515645744 7288x^{56} - 221382136587677 4435x^{57} \\ &- 260822367118104 1657x^{58} + 196280141142053 07016x^{59} + 651404631123094716x^{60} \\ &+ 761528771717834208x^{61} + 888518897148324570x^{62} + 1034700084660340082x^{63} \\ &- 752584720886834 3576x^{64} - 116633198562800052x^{65} - 134898641118903336x^{66} \\ &- 155761841308153260x^{67} - 179556106888260384x^{68} + 1266995051549992032x^{69}\end{aligned}$$

$$
\begin{aligned}
g_{24}(x) = {} & 44199864566676 + 53277891868890x + 36951353492365x^2 - 18447292166275x^3 \\
& - 1159818177616 56x^4 + 5683172059809972x^5 + 7671280571862570x^6 \\
& + 6327550580716885x^7 - 1472857823809435x^8 - 18209145388579992x^9 \\
& + 496676536774208388x^{10} + 771501576163548690x^{11} + 774677809879397185x^{12} \\
& + 37151453118769745x^{13} - 2080007375935924008x^{14} + 449042986342764651 24x^{15} \\
& + 851924419367526376 50x^{16} + 111451370786904498265x^{17} + 478091161409322558 65x^{18} \\
& - 289357227498865856904x^{19} + 6805964762978995053180x^{20} + 19701780140886237080850x^{21} \\
& + 479205560785200208558 75x^{22} + 10533749951691408844 3775x^{23} \\
& - 176316318895755129455 8680x^{24} - 16821499115216128484 40924x^{25} \\
& - 31440911069600555059 26510x^{26} - 56825607629261748192 91685x^{27} \\
& - 99831139953378645033 02425x^{28} + 14584420902969065396 6024044x^{29} \\
& + 90577048940932553337666992x^{30} + 14901633736857070123 6013970x^{31} \\
& + 24098797717913362853 8814635x^{32} + 38370773386942593957 3259415x^{33} \\
& - 50546371860993653129 20130012x^{34} - 23830735426363507644 33826072x^{35} \\
& - 36462478787448597914 70683310x^{36} - 55169276506090399184 96642165x^{37} \\
& - 82607620583110105972 10260105x^{38} + 99947418327478671197 343833527x^{39} \\
& + 37844995423440695266 939743959x^{40} + 55100153059398963201 126006300x^{41} \\
& + 79573062038413273868 028870415x^{42} + 11403486438583866058 9245973115x^{43} \\
& - 12838559200275215856 01121843789x^{44} - 40163143048418250301 1941004665x^{45} \\
& - 56341855962152203891 5547815300x^{46} - 78542236109552227527 3631000625x^{47} \\
& - 10883436566066677478 39725560325x^{48} + 11515350760355635501 320142255915x^{49} \\
& + 30239361383197553614 42261262591x^{50} + 41194126526500884037 66862240940x^{51} \\
& + 55833989738214014374 27782781615x^{52} + 75309075393777920432 31905759675x^{53} \\
& - 75471967365836479567 646155794821x^{54} - 16787688185166611656 225020168433x^{55} \\
& - 22327203688191441773 802872732580x^{56} - 29570337359172614836 200572834465x^{57} \\
& - 39004838361833906826 391115026885x^{58} + 37258730284260242337 4479932324863x^{59} \\
& + 70520166819123145016 529674257763x^{60} + 91921490099273043258 578272890040x^{61} \\
& + 11939341747619524428 640880176663 5x^{62} + 15454327717036080214 6843655221595x^{63} \\
& - 14144604509430612129 79844779136033x^{64} - 22805858627534448604 0909532074621x^{65} \\
& - 29219816324086943703 6528542283800x^{66} - 37323699985993013918 1490117490085x^{67} \\
& - 47534021904105270921 4076298517285x^{68} + 41866419213874631136 360904278657 91x^{69} \\
& + 57398239959744024110 4412076252915x^{70} + 72450880852088947486 9872176906800x^{71} \\
& + 91208712819458218060 6507235839875x^{72} + 11452641819362424724 3304052045807 5x^{73} \\
& - 97428738783150914112 50160134457665x^{74} - 11304132801034427539 36876665890989x^{75} \\
& - 14082643648408458310 4413943778 4160x^{76} - 17503195352562682381 32497892063885x^{77}
\end{aligned}
$$

$$- 217050292292597450614356666109072 5x^{78} + 178908346396081800128050542896422 59x^{79}$$
$$+ 174273927157410827127379380571162 7x^{80} + 214596668243174027346943594416359 5x^{81}$$
$$+ 263702290720010115399708762459186 0x^{82} + 323388616454863960514654032186951 5x^{83}$$
$$- 258979696654933874453663498838365 97x^{84} - 209277041236551323843525466560523 7x^{85}$$
$$- 255027677869883451948277441423428 5x^{86} - 310205015395566557899404046596114 0x^{87}$$
$$- 376636488968328711677534384551060 5x^{88} + 293740000624483804478890735475612 67x^{89}$$
$$+ 193524003381979598103941509210061 9x^{90} + 233626540368501650513297119546997 5x^{91}$$
$$+ 281569761954187989276744030598524 0x^{92} + 338797433941009471781532595632301 5x^{93}$$
$$- 257859241059525699489280041123788 49x^{94} - 135043158279578220375813666741080 5x^{95}$$
$$- 161644507552087739540574171296922 5x^{96} - 193194550745668914394263466061700 0x^{97}$$
$$- 230561116935915600353154200552902 5x^{98} + 171564186963047636005504085621510 55x^{99}$$
$$+ 687379082077441716165690906155771 x^{100} + 816415958977468475853193747835215 x^{101}$$
$$+ 968348850584524348447504741362840 x^{102} + 114701061071380775903904309509662 5x^{103}$$
$$- 835819515053050842041607702170045 1x^{104} - 240714899339982433449408005104533 x^{105}$$
$$- 283875024040741105674985093276305 x^{106} - 334355514938579365555961289992040 x^{107}$$
$$- 393328645903551948398787580245735 x^{108} + 281083558081591296236258923424361 3x^{109}$$
$$+ 518278329726246724811560651432 83x^{110} + 607215335910078944325739669476 15x^{111}$$
$$+ 710599724686687255376595889863 60x^{112} + 830648912806717372603328438262 45x^{113}$$
$$- 582904099186926848986624808653503 x^{114} - 517019332694789181790852074758 1x^{115}$$
$$- 602086116545798442084371964262 5x^{116} - 700411474038816702928561394196 0x^{117}$$
$$- 813949160024358721240736811283 5x^{118} + 561545826786078371225961013512 51x^{119}$$

References

[1] X. Hou, A class of permutation binomials over finite fields, J Number Theory 133 (2013), 3549-3558

[2] X. Hou, Determination of a type of permutation trinomials over finite fields, II, arXiv:1404.1822, 2014

[3] X. Hou and S. D. Lappano, Determination of a type of permutation binomials over finite fields, arXiv:1312.5283, 2013

[4] R. Lidl and H. Niederreiter, Finite Fields, @nd ed., Cambridge Univ. Press, Cambridge, 1997

[5] M. E. Zieve, *Permutation Polynomials on $\mathbb{F}_q$ induced from Redei function bijections on subgroups of $\mathbb{F}_q^*$, arXiv:1310.0776,2013*

Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620

E-mail address: slappano@mail.usf.edu